

INTEGRAL CLOSURES OF COHEN-MACAULAY MONOMIAL IDEALS

ABDUL SALAM JARRAH

ABSTRACT. The purpose of this paper is to present a family of Cohen-Macaulay monomial ideals such that their integral closures have embedded components and hence are not Cohen-Macaulay.

1. INTRODUCTION

Let R be an arbitrary ring. Krull [5] asked whether there exists a primary ideal I such that the integral closure $\bar{I}$ of I has embedded primes. Huneke [4] gave a counterexample, his example in a 3-dimensional regular local ring of characteristic 2. No other examples have appeared in the literature. The long history of Krull's question and the machinery that is required to understand Huneke's example suggest that there might not be an easy counterexample. In fact, it is easy to see that if I is a P -primary monomial ideal in a polynomial ring, then the integral closure $\bar{I}$ of I is a P -primary monomial ideal. Thus, there is no monomial ideal that can serve as a counterexample to Krull's question. Vasconcelos asked a relaxed version of Krull's question, namely whether there exists an unmixed monomial ideal I in a polynomial ring such that the integral closure $\bar{I}$ has embedded primes.

In this paper, I present a family of Cohen-Macaulay monomial ideals such that their integral closures are not Cohen-Macaulay, in particular, their integral closures have embedded primes. Thus, each example in this family is a counterexample to Vasconcelos's question. Naturally, this family is not the only family of monomial ideals with this property. Another Cohen-Macaulay monomial ideal is given at the end which does not belong to the family, but whose integral closure has an embedded prime.

Throughout this paper, n and t are positive integers and $n \geq 3$.

2. EXAMPLES

Let $R = k[x_1, \dots, x_n] = k[\mathbf{x}]$ be a polynomial ring over an arbitrary field k . For any positive integer m , let $[m] = \{1, \dots, m\}$. For $i \in [n]$, let $e_i = (t, \dots, t, 0, t, \dots, t) \in \mathbb{R}^n$, with 0 is in the i -th coordinate. Thus the corresponding monomial with exponent e_i is $\mathbf{x}^{e_i} = x_1^t \cdots x_{i-1}^t x_{i+1}^t \cdots x_n^t$. Consider the monomial ideal

$$I_{n,t} = \langle \mathbf{x}^{e_1}, \dots, \mathbf{x}^{e_n} \rangle.$$

Date: December 4, 2001.

For $i, j \in [n]$ and $i \neq j$, it is easy to see that $I_{n,t} \subset \langle x_i^t, x_j^t \rangle$. Thus $I_{n,t} \subseteq \bigcap_{i=1}^{n-1} \bigcap_{j=i+1}^n \langle x_i^t, x_j^t \rangle$ and it is easy to see that equality holds. Moreover, without loss of generality, $\bigcap_{i=2}^{n-1} \bigcap_{j=i+1}^n \langle x_i^t, x_j^t \rangle \not\subseteq \langle x_1^t, x_2^t \rangle$, since $x_3^t \cdots x_n^t \in \bigcap_{i=2}^{n-1} \bigcap_{j=i+1}^n \langle x_i^t, x_j^t \rangle$ but $x_3^t \cdots x_n^t \notin \langle x_1^t, x_2^t \rangle$. Therefore,

$$I_{n,t} = \bigcap_{i=1}^{n-1} \bigcap_{j=i+1}^n \langle x_i^t, x_j^t \rangle$$

is the minimal primary decomposition of $I_{n,t}$. In particular, $I_{n,t}$ has no embedded primes and $\text{codim}(I_{n,t}) = 2$. On the other hand, the minimal free resolution of $R/I_{n,t}$ is

$$0 \leftarrow R/I_{n,t} \leftarrow R \xleftarrow{\begin{pmatrix} \mathbf{x}^{e_1} & \cdots & \mathbf{x}^{e_n} \end{pmatrix}} R^n \xleftarrow{\begin{pmatrix} -x_1^t & -x_1^t & \cdots & -x_1^t \\ x_2^t & 0 & \cdots & 0 \\ 0 & x_3^t & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & x_n^t \end{pmatrix}} R^{n-1} \leftarrow 0.$$

Hence, the projective dimension of $R/I_{n,t}$ equals 2. Since $\text{codim}(I_{n,t})$ is equal to the projective dimension of $R/I_{n,t}$, $I_{n,t}$ is Cohen-Macaulay.

Let Γ be the set of exponents of monomials of $I_{n,t}$. Let Ω be the convex hull of Γ in $\mathbb{R}_{\geq 0}^n$.

Theorem 2.1. *Let $(a_1, \dots, a_n) \in \Omega$. Then, for $s \in [n]$ and $\{i_1, \dots, i_s\} \subseteq [n]$, we have*

$$a_{i_1} + \cdots + a_{i_s} \geq t(s-1).$$

Proof. Since Ω is the convex hull of exponents of monomials of $I_{n,t}$ in $\mathbb{R}_{\geq 0}^n$ and $I_{n,t} = \langle \mathbf{x}^{e_1}, \dots, \mathbf{x}^{e_n} \rangle$, there exists, for $i \in [n]$, $c_i \geq 0$ and $d_i \geq 0$ such that $c_1 + \cdots + c_n = 1$ and

$$(a_1, \dots, a_j, \dots, a_n) = c_1 e_1 + \cdots + c_n e_n + (d_1, \dots, d_n).$$

Thus

$$\begin{aligned} (a_1, \dots, a_j, \dots, a_n) &\geq c_1 e_1 + \cdots + c_n e_n \\ &= (t \sum_{i=2}^n c_i, \dots, t \sum_{\substack{i=1 \\ i \neq j}}^n c_i, \dots, t \sum_{i=1}^{n-1} c_i). \end{aligned}$$

In particular, for $j \in [n]$, we have $a_j \geq t \sum_{\substack{i=1 \\ i \neq j}}^n c_i$. Therefore, for $\{i_1, \dots, i_s\} \subseteq [n]$, we have

$$\begin{aligned} a_{i_1} + \dots + a_{i_s} &\geq t \left(\sum_{\substack{i=1 \\ i \neq i_1}}^n c_i + \dots + \sum_{\substack{i=1 \\ i \neq i_s}}^n c_i \right) \\ &= t(-c_{i_1} - \dots - c_{i_s} + s \sum_{i=1}^n c_i) \\ &\geq t(s-1). \end{aligned}$$

□

It is well-known, see [3, Exerice 4.23], that the integral closure $\bar{I}_{n,t}$ of $I_{n,t}$ is the monomial ideal generated by the integral points in Ω . Thus

$$\bar{I}_{n,t} = \langle x_1^{a_1} \dots x_n^{a_n} : (a_1, \dots, a_n) \in \Omega \cap (\mathbb{N} \cup \{0\})^n \rangle.$$

Let

$$\begin{aligned} \Delta_{n,t} := & \{(a_1, \dots, a_n) : a_i \in [t] \text{ for all } i \text{ and } a_1 + \dots + a_n = t(n-1)\} \\ & \cup \{e_i : i = 1, \dots, n\}. \end{aligned}$$

The following theorem gives a finite set of generators of $\bar{I}_{n,t}$.

Theorem 2.2. *Let $\Delta_{n,t}$ and $I_{n,t}$ be as above. Then*

$$\bar{I}_{n,t} = \langle x_1^{a_1} \dots x_n^{a_n} : (a_1, \dots, a_n) \in \Delta_{n,t} \rangle.$$

Proof. Let $(a_1, \dots, a_n) \in \Delta_{n,t}$. For $j \in [n]$, let

$$c_j = 1 - \frac{a_j}{t}.$$

It is straightforward to check that $c_j \geq 0$, $c_1 + \dots + c_n = 1$, and $c_1 e_1 + \dots + c_n e_n = (a_1, \dots, a_n)$. Therefore, $(a_1, \dots, a_n)$ is a convex combination of $e_1, \dots, e_n$. Thus $(a_1, \dots, a_n) \in \Omega$, and hence $x_1^{a_1} \dots x_n^{a_n} \in \bar{I}_{n,t}$.

To prove the other inclusion, let $(b_1, \dots, b_n) \in \Omega \cap (\mathbb{N} \cup \{0\})^n$. It is enough to find $(a_1, \dots, a_n) \in \Delta_{n,t}$ such that $a_i \leq b_i$ for all i . But $I_{n,t} \subseteq \langle x_1^{a_1} \dots x_n^{a_n} : (a_1, \dots, a_n) \in \Delta_{n,t} \rangle$, so we need to prove the above statement only for the case when $x_1^{b_1} \dots x_n^{b_n} \in \bar{I}_{n,t} \setminus I_{n,t}$. Since $x_1^{b_1} \dots x_n^{b_n} \notin I_{n,t}$, there exists $s \geq 2$ and $\{i_1, \dots, i_s\} \subset [t]$ such that $b_{i_1}, \dots, b_{i_s} < t$. Let s be maximal with this property and without lose of generality, assume $\{i_1, \dots, i_s\} = \{1, \dots, s\}$. Then $b_1, \dots, b_s < t$ and $b_{s+1}, \dots, b_n \geq t$. Let $a_{s+1} = ts - (b_1 + \dots + b_s)$. By Theorem 2.1, $a_{s+1} \leq ts - t(s-1) = t$. Also, $a_{s+1} > 0$, since $b_{i_1}, \dots, b_{i_s} < t$. Thus $a_{s+1} \in [t]$. It is clear that the n -tuple $(b_1, \dots, b_s, a_{s+1}, t, \dots, t) \in \Delta_{n,t}$ and $x_1^{b_1} \dots x_s^{b_s} x_{s+1}^{a_{s+1}} x_{s+1}^t \dots x_n^t$ divides $x_1^{b_1} \dots x_n^{b_n}$. We conclude that $x_1^{b_1} \dots x_n^{b_n} \in \langle x_1^{a_1} \dots x_n^{a_n} : (a_1, \dots, a_n) \in \Delta_{n,t} \rangle$. □

Example 2.3. In Figure 1, the vertices of the triangle correspond to the generators of the ideal $I_{3,4}$. Also, all the dots inside and on the boundary of the triangle correspond to the generators of the integral closure $\bar{I}_{3,4}$ of $I_{3,4}$.




FIGURE 1. $I_{3,4}$, its convex hull, and the generators of $\bar{I}_{3,4}$.

The following theorem shows that $\bar{I}_{n,t}$ has an embedded associated prime for $t \geq 2$.

Theorem 2.4. *For $t \geq 2$, the integral closure $\bar{I}_{n,t}$ has $\langle x_1, x_2, x_3 \rangle$ as an embedded associated prime.*

Proof. By Theorem 2.2, it is easy to see that $(\bar{I}_{n,t} : x_1 x_2^{t-1} x_3^{t-1} x_4^t \cdots x_n^t) = \langle x_1, x_2, x_3 \rangle$. Hence, by [1, Theorem 4.5], $\langle x_1, x_2, x_3 \rangle \in \text{Ass}(R/\bar{I}_{n,t})$. Using the same argument, we find that $(\bar{I}_{n,t} : x_2^{t-1} x_3^t \cdots x_n^t) = \langle x_1, x_2 \rangle$ and hence $\langle x_1, x_2 \rangle \in \text{Ass}(R/\bar{I}_{n,t})$. Thus $\langle x_1, x_2, x_3 \rangle$ is an embedded prime. $\square$

Analogously, $\bar{I}_{n,t}$ has at least $\binom{n}{3}$ embedded primes, one for each triple of variables. Since $\bar{I}_{n,t}$ has embedded primes, by [6, Theorem 17.3], the following corollary is straightforward.

Corollary 2.5. *The ideal $\bar{I}_{n,t}$ is not Cohen-Macaulay for all $t \geq 2$.*

In the following example, we compute the ideal $\bar{I}_{3,2}$.

Example 2.6. $I_{3,2} = \langle x_1^2 x_2^2, x_1^2 x_3^2, x_2^2 x_3^2 \rangle = \langle x_2^2, x_3^2 \rangle \cap \langle x_1^2, x_3^2 \rangle \cap \langle x_1^2, x_2^2 \rangle$. By Theorem 2.2, it is easy to see that

$$\bar{I}_{3,2} = \langle x_1^2 x_2^2, x_1^2 x_3^2, x_2^2 x_3^2, x_1 x_2 x_3^2, x_1 x_2^2 x_3, x_1^2 x_2 x_3 \rangle.$$

Moreover,

$$\bar{I}_{3,2} = \langle x_2^2, x_2x_3, x_3^2 \rangle \cap \langle x_1^2, x_1x_3, x_3^2 \rangle \cap \langle x_1^2, x_1x_2, x_2^2 \rangle \cap \langle x_1^2, x_2^2, x_3^2 \rangle.$$

It is clear that $\langle x_1^2, x_2^2, x_3^2 \rangle$ is an embedded component of $\bar{I}_{3,2}$.

I have found many other families of examples but I presented only the simplest one. In the following example, I give a Cohen-Macaulay monomial ideal which is not in the family above and whose integral closure is not Cohen-Macaulay.

Example 2.7. Let $S = k[x, y, z, w]$ be a polynomial ring over an arbitrary field k . Let $I = \langle x^3yz, x^2w^2, y^2w^3 \rangle = \langle x^2, y^2 \rangle \cap \langle x^3, x^2w^2, w^3 \rangle \cap \langle y, w^2 \rangle \cap \langle z, w^2 \rangle$. Since all the components have codimension 2, there are no inclusion relations among them, so by the uniqueness of minimal primary components, the above decomposition of I is irredundant. Thus I is a generic (see [7, Definition 1.1]) monomial ideal with no embedded primes. By [7, Theorem 2.5], I is Cohen-Macaulay. Now by using the computer algebra system *Normaliz* [2], the integral closure $\bar{I}$ of I is

$$\bar{I} = \langle x^3yz, x^2w^2, y^2w^3, x^2y^2zw, xy^2zw^2, xyw^3 \rangle.$$

It is easy to see that $(\bar{I} : xyw^2) = \langle x, w \rangle$ and $(\bar{I} : xy^2w^2) = \langle x, z, w \rangle$. Thus $\bar{I}$ has $\langle x, z, w \rangle$ as an embedded prime and hence is not Cohen-Macaulay.

3. ACKNOWLEDGMENTS

I thank Professor Irena Swanson for suggesting the problem, and for helpful discussions and conversations. Also, I thank the referee for pointing out errors in the earlier version of this paper.

REFERENCES

- [1] M. Atiyah and I. Macdonald, *Introduction to commutative algebra*, Addison-Wesley Publishing Company, 1969.
- [2] W. Bruns and R. Koch, *NORMALIZ*, a program for computing normalizations of affine semigroups (1998). Available by anonymous ftp from <ftp://ftp.mathematik.uni-osnabrueck.de/pub/osm/kommalg/software/>.
- [3] D. Eisenbud, *Commutative algebra with a view toward Algebraic Geometry*, Graduate Texts in Mathematics **150**, Springer, 1996.
- [4] C. Huneke, The primary components of and integral closures of ideals in 3-dimensional regular local rings, *Math. Ann.* **275** (1986), 617–635.
- [5] W. Krull, Beiträge zur Arithmetik kommutativer Integritätsbereiche, *Math. Z.* **41** (1936), 545–577.
- [6] H. Matsumura, *Commutative ring theory*, Cambridge Studies in Advanced Mathematics **8**, 1986.
- [7] E. Miller, B. Sturmfels, and K. Yanagawa, Generic and cogenerated monomial ideals, *J. Symbolic Comp.* **29** (2000), 691–708.

(Abdul Salam Jarrah) DEPARTMENT OF MATHEMATICAL SCIENCES, NEW MEXICO STATE UNIVERSITY, LAS CRUCES, NM 88003, USA

E-mail address: ajarrah@nmsu.edu